

## A rare case of amino nitrogen poisoning with methemoglobinemia and severe metabolic acidosis

Y Sai Vani<sup>1\*</sup>, M Suma Pravalika<sup>2</sup>

<sup>1</sup>HOD, <sup>2</sup>Clinical Pharmacist, Dept. of Nephrology, Santhiram Medical College, Nandyala, Andhra Pradesh, India

**\*Corresponding Author: Y Sai Vani**

Email: drsaivani@gmail.com

---

### Abstract

Methemoglobinemia can be managed medically with methylene blue but in severe methemoglobinemia sometimes extracorporeal therapy may be required, here we report a rare case of Amino nitrogen poisoning who presented with severe metabolic acidosis, shock and had severe methemoglobinemia that was treated with methylene blue and haemo dialysis.

**Keywords:** Amino, Methemoglobinemia.

---

### Introduction

Amino nitrogen poisoning is rare and methemoglobinemia due to this compound is not yet reported, the nitrogen compound is converted into nitrites by bacteria in our saliva, stomach, and intestines, and it is primarily the nitrites that cause toxicity. Nitrites oxidize the iron component of red blood cells (haemoglobin), rendering them unable to carry oxygen. The resulting condition is called methemoglobinemia, and the lack of oxygen is the reason behind the characteristic pale to blue-gray color of the skin. The severity of methemoglobinemia is directly proportional to the percentage of red blood cells affected. The higher the percentage, the more serious the symptoms, and the bluer the patient appears.

Ferrous iron can be converted to ferric iron by oxidative stress which results in the formation of methemoglobin.<sup>1</sup> The iron moiety in hemoglobin exists in the “ferrous” state and undergoes oxidation to the “ferric” state forming methemoglobin at a rate of ~3% per day.<sup>2</sup> The initial sign of methemoglobinemia is cyanosis<sup>3</sup> and the diagnosis should be considered in all patients who present with cyanosis, particularly if it does not improve with supplemental oxygen. Methemoglobin levels of >70% are generally lethal, although survival has been reported with a level of 94%.<sup>4</sup> Methylene blue is indicated as the first-line antidote therapy for patients with severe methemoglobinemia.<sup>5</sup> Patients with methemoglobinemia typically present with skin discoloration (“chocolate cyanosis”), especially of the

nails, lips, and ears, whereas the color is more often brown rather than blue. Even at low levels of MetHb, this discoloration can be striking.<sup>6</sup> Methylene blue should be used with caution in individuals with severe renal insufficiency and in young patients with G6PD deficiency. This group of patients can develop hemolytic anemia characterized by formation of Heinz bodies without any reduction in MetHb levels.<sup>6-8</sup> Methemoglobin forms when the ferrous (Fe<sup>2+</sup>) form of heme in hemoglobin is oxidized to the ferric (Fe<sup>3+</sup>) state. Ferric heme is unable to bind oxygen, and the remaining ferrous heme develops increased oxygen affinity and causes impaired oxygen delivery to tissues, resulting in a “left shift” in the oxygen dissociation curve.<sup>9</sup>

Which results in hypoxic ischemia to the tissues and causes severe lactic acidosis, now we report a case of amino nitrogen poisoning with severe methemoglobinemia and severe metabolic acidosis.

### Case Report

A previously healthy, 34-year male came with the chief complaints of self-ingestion of Amino nitrogen of 350ml, at his residence at 6:30pm. Gastric lavage was performed at the local hospital and came to SRMC at 8:30 pm for further management. On his arrival the patient was in gasping state and CPR was done for cardiac arrest. Patient was intubated in the emergency ward. Later the patient was on mechanical ventilation. On examination, he was afebrile (98.6°F), with heart

rate 120/min, regular BP 116/72, respiratory rate was 20/min and the oxygen saturation is 80% of room air latter he was becoming hypotensive and inotropes were initiated.

Presenting symptoms of the patient were

1. Bluish skin due to lack of oxygen (noticed around mouth/lips first)
2. Shortness of breath
3. Nausea, vomiting
4. Dehydration (from loss of bodily fluids)
5. Tachycardia



**Fig. 1:**

Patient was shifted to critical care medicine (CCM) for further management and observation. In CCM, he received inotropes like Noradrenaline and Vasopressin to maintain his vitals which was shown in the Table 1. The patient was cyanosed had both central and peripheral cyanosis with hypoxia and had metabolic acidosis with PH of 7.052, so diagnosis of Amino

nitrogen ingestion with severe central and peripheral cyanosis, (Fig.1) methemoglobinemia was suspected and Methylene blue was initiated, the patient had severe metabolic acidosis, so dialysis (Hemo diafiltration) was initiated with blood flow of 150, dialysate flow of 300, heparin free, Ultra filtration nil and Methylene blue injection @ 1mg/kg (100mg) IV was given over 20 minutes. After initiation of methylene blue and Hemodialysis, inotropic support was decreased and was stopped after 24hrs. His haemoglobin levels were 15.2g/dl and methaemoglobin levels were 60%. After 2 days the methaemoglobin levels were decreased to 58%.

Initial Renal function tests, Arterial Blood Gas were enclosed in tables. The patient was maintained on high Po<sub>2</sub> (hyper baric oxygen) to improve oxygenation initially by increasing the dissolved oxygen and oxygen saturations in pulse oximeter was not considered because after giving Methylene blue they were not significant, we were depending on lactates in ABG (Arterial Blood Gas analysis) shown in the (table 2), to know the perfusion of tissues and oxygenation daily depending on them we were maintaining the paO<sub>2</sub> by altering the FI<sub>O2</sub> and ventilator modes, the ventilator parameters were shown in the Table 4.

The molecular weight of amino nitrogen is 29 g/mol and it is water soluble excreted through kidneys even then we have dialysed him 3 times as his methemoglobin levels were high for first 4 days in order to decrease the poison load from the body though his metabolic parameters were corrected and Renal Function Tests (RFTs) were normal, and he was extubated on 7<sup>th</sup> day, he had DVT due to femoral catheter and started on 5000u heparin twice a day. He was discharged on 14<sup>th</sup> day of his admission after subsidence of DVT.

**Table 1:**

On arrival at ER	Pulse	BP	Spo2	Inotropics	
				Noradrenaline	Adrenaline
6 hrs	132	118/77	79	10ml	0.5ml
12 hrs	128	139/73	99	4ml	0.5ml
1 <sup>st</sup> day	132	118/77	79	NO	NO
2 <sup>nd</sup> day	100	168/75	76	NO	NO
3 <sup>rd</sup> day	113	130/81	83	NO	NO

4 <sup>th</sup> day	103	140/86	78	NO	NO
5 <sup>th</sup> day	183	154/88	100	NO	NO
6 <sup>th</sup> day	132	118/78	100	NO	NO
7 <sup>th</sup> day	118	141/65	67	NO	NO

**Table 2:** ABG values

Date& Time	Blood Gas values			Electrolyte & Metabolite values					Oximetry values	Calculated values	
	pH	pCO <sub>2</sub>	pO <sub>2</sub>	cK <sup>+</sup>	cNa <sup>+</sup>	cCa <sup>2+</sup>	cCl <sup>-</sup>	cLac	ctHb	sO <sub>2</sub>	Hct <sub>c</sub>
11/10&3:51AM	7.052	30.2	366	3.9	149	1.03	114	19	16.3	-	49.9
11/9&8:28AM	7.465	29.1	107	2.5	16	0.60	128	0.9	4.5	98.2	14.4
11/10&7:31AM	7.267	38.2	188	4.0	147	1.07	114	8.2	17.2	97.1	52.6
11/10&6:31PM	7.423	39.4	233	3.6	139	1.07	111	10.4	14.7	97.0	45.1
11/10&7:30PM	7.374	39.1	480	3.4	141	1.06	111	11.5	14.8	97.6	45.3
11/11&6:42AM	7.468	37.7	491	3.5	141	1.06	112	10.8	13.3	96.8	40.9
	7.533	28.6	465	3.4	140	1.09	111	12.4	13.0	96.8	39.8
11/12&6:38AM	7.492	31.5	332	3.3	139	1.08	112	10.4	12.5	96.0	38.4

**Table 3:** Dialysis chart

S. No	Day from admission	QB	QD	Heparin	Ultra filtration	Duration	Pulse rate	Blood Pressure	sPO <sub>2</sub>	HCO <sub>3</sub> <sup>-</sup>
1	Day 1	150	300	free	Nil	3hrs	86	107/70	80	8.4
2	Day 3	150	300	free	Nil	6hrs	82	120/80	76	26.3
3	Day 4	150	300	free	Nil	6hrs	114	145/82	98	27.3

**Table 4:** Ventilator settings

Date	Mode of ventilation	FiO <sub>2</sub>	Respiratory Rate	Tidal Volume	PEEP
10/11/2020	AC/PC	110	20	599	8
11/11/2020	AC/PC	90	20	372	10
12/11/2020	ADL/PCV	80	25	301	10
13/11/2020	ADL/PCV	75	21	550	10
14/11/2020	PCV/CPAP	100	27	553	10
15/11/2020	CPAP	35	37	480	9.8

**Table 5:** Renal function tests

Date	Urea	Creatinine	Uric Acid
10/11/2020	20	1.35	8.2
11/11/2020	38	0.60	5.6
12/11/2020	43	0.84	3.2
13/11/2020	42	0.82	5.9
14/11/2020	39	0.70	3.4
15/11/2020	39	0.55	3.5

## Discussion

Methylene blue is used to help make oxygen available to the body by reversing what nitrates have done to the molecules that carry oxygen in the body. His haemoglobin levels were 15.2g/dl and methaemoglobin levels were 60%. After 2 days the methaemoglobin levels were decreased to 58%. Methemoglobin levels of >70% are generally lethal, although survival has been reported with a level of 94%.<sup>10</sup> In 1959, aniline poisoning has been first described in a case series of marking-ink poisoning causing methemoglobinemia in babies.<sup>11</sup> Later, more cases have been published, most often in correlation with dye or herbicides exposure.<sup>12-15</sup> Hyperbaric oxygen, either alone or in combination with methylene blue, has been successfully used for management of methemoglobinemia.<sup>16-18</sup> A case of methemoglobinemia occurred after the application of silver nitrate solution to burns; the production of nitrite was shown to be due to *Aerobacter cloacae* present on the skin.<sup>19</sup> Three patients who were burned with molten sodium and potassium nitrate in an explosion developed methemoglobinemia and one patient died.<sup>20</sup> However, eight cases of severe methemoglobinemia were reported in patients with some degree of renal impairment or gastrointestinal dysfunction, following the ingestion of therapeutic doses of ammonium nitrate over periods ranging from 2 to 46 days.<sup>21-23</sup> Methylene blue should be used with caution in individuals with severe renal insufficiency and in young patients with G6PD deficiency. This group of patients can develop hemolytic anemia characterized by formation of Heinz bodies without any reduction in MetHb levels.<sup>6-8</sup> Alternative therapies include exchange transfusion, hyperbaric oxygen and ascorbic acid.<sup>24</sup>

## Conclusion

The patient had Central and peripheral cyanosis and severe metabolic acidosis, due to amino nitrogen poisoning, which is not yet reported, he required not only medical management for poisoning with methylene blue but also Haemo dialysis in view of severe metabolic acidosis and after to support the kidneys in excretion of residual poison in view of severe methemoglobinemia, till the crisis period the person was maintained on hyperbaric oxygen to improve

oxygenation of the tissues and regular lactates were monitored for clinical decisions.

## Aniline

## Conflict of Interest

None.

## Sources of Funding

None.

## References

1. Shatila W, Verma A, Adam S. Plasmapheresis in severe methemoglobinemia following occupational exposure. *Transfus Apher Sci*. 2017 Jun;56(3):341-344. doi: 10.1016/j.transci.2017.02.003.
2. Skold A, Cosco DL, Klein R. Methemoglobinemia: pathogenesis, diagnosis, and management. *South Med J* 2011;104(11):757-61.
3. CM Alexander, LE Teller, Gross JB, "Principles of pulse oximetry: theoretical and practical considerations," *Anesth Analgesia* 1989;68(3):368-76.
4. Harvey M, Cave G, Chanwai G, "Fatal methaemoglobinaemia induced by self-poisoning with sodium nitrite," *Emerg Med Aust* 2010;22(5):463-5.
5. Umbreit J, "Methemoglobin—it's not just blue: a concise review," *Am J Hematol* 2007;82(2):134-44.
6. Olsen KR, "Poisoning and drug overdose," in *Methylene Blue by Fabian Garza*, pp. 902-904, McGraw-Hill, New York, NY, USA, 5th edition, 2007.
7. Clifton II J, Leikin JB, "Methylene blue," *Am J Ther* 2003;10(4):289-91.
8. Sikka P, Bindra VK, Kapoor S, Jain V, Saxena KK, "Blue cures blue but be cautious," *J Pharm Bioallied Sci*, 2011;3(4):543-5.
9. Darling R, Roughton F. Effect of methemoglobin on the equilibrium between oxygen and hemoglobin. *Am J Physiol* 1942;137:56.
10. Harvey M, Cave G, Chanwai G, "Fatal methaemoglobinaemia induced by self-poisoning with sodium nitrite," *Emerg Med Aust* 2010;22(5):463-5.
11. Ramsay DE, Harvey C, "Marking-ink poisoning: an outbreak of methemoglobin cyanosis in newborn babies," *Lancet* 1959;273(7079):910-2.
12. Pizon AF, Schwartz AR, Shum LM. "Toxicology laboratory analysis and human exposure to p-chloroaniline," *Clin Toxicol* 2009;47(2):132-6.
13. Bazylewicz A, Kłopotowski T, Kicka M, Miśkiewicz Ł, Picheta S, "Acute poisoning due to chemical substances inducing methemoglobinemia—two cases report," *Przegląd Lekarski* 2010;67(8):636-9.
14. Watt BE, Proudfoot AT, Bradberry SM, Vale JA, "Poisoning due to urea herbicides," *Toxicol Rev* 2005;24(3):161-6.
15. Thier R, Lewalter J, Selinski S, Bolt HM, "Biological monitoring in workers in a nitrobenzene reduction

- plant:haemoglobin versus serum albumin adducts," *Int Arch Occup Environ Health* 2001;74(7):483-8.
16. Goldstein GM, Doull J. Treatment of nitrite-induced methemoglobinemia with hyperbaric oxygen. *Proc Soc Exp Biol Med* 1971;138(1):137-9.
  17. Goulon M, Nouailhat F, Gajdos P. On a case of acquired methemoglobinemia with coma, treated with hyperbaric oxygen and methylene blue. *Rev Neurol (Paris)* 1966;114(5):376-8.
  18. Jansen T, Barnung S, Mortensen CR, Jansen EC. Isobutyl-nitrite-induced methemoglobinemia; treatment with an exchange blood transfusion during hyperbaric oxygenation. *Acta Anaesthesiol Scand* 2003;47(10):1300-1.
  19. Ternberg JL, Lute E: Methemoglobinemia: a complication of the silver nitrate treatment of burns. *Ped Surg* 1968;63:328-30.
  20. Harris JC, Rumack BH, Peterson RG. Methemoglobinemia resulting from absorption of nitrates. *JAMA* 1979;242:2869-71.
  21. Keith NM, Whelan M, Bannick EG: The action and excretion of nitrates. *Arch Znt h4ed* 1930;46:797-832.
  22. Eusterman GB, Keith NM: Transient methemoglobinemia following the administration of ammonium nitrate. *A4ed Clin North Am* 1926;12:1489-96.
  23. Tarr L. Transient methemoglobinemia due to ammonium nitrate. *Arch Znt Med* 1933;51:38.
  24. Boran P, Tokuc G, Yegin Z, "Methemoglobinemia due to application of prilocaine during circumcision and the effect of ascorbic acid," *J Pediatric Urol* 2008;4(6):475-6.

**How to cite:** Vani YS, Pravalika MS. A rare case of amino nitrogen poisoning with methemoglobinemia and severe metabolic acidosis. *IP J Urol Nephrol Hepatol Sci* 2020;3(4):92-6.