

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Comprehensive and Advanced Pharmacology

Journal homepage: <https://www.ijcap.in/>

Original Research Article

A chemical rationale of drug stability and degradation- An insightful approach

Senthilraja. M^{1,*}

¹Dept. of Pharmaceutical Chemistry, Kasthooribha Gandhi Pharmacy College, Rasipuram, Tamil Nadu, India



ARTICLE INFO

Article history:

Received 20-11-2021

Accepted 21-04-2022

Available online 16-05-2022

Keywords:

Drug

Stability

Degradation

Hydrolysis

Prodrug Strategy

ABSTRACT

To discuss the chemical rationale for the stability of certain functional groups within drug molecules against reaction such as hydrolytic degradation and consider the methods used to maximize their stability on storage.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The metabolism of drugs occurs through basic chemical reactions as soon as the administered compound comes into contact with enzymes that are capable of altering its chemical structure. Conversely, a drug's stability after administration is due mainly to its lack of transformation by the body's enzymes. However, many drugs are susceptible to some form of chemical decomposition, be it through the interaction with enzymes or through improper storage and use, and such degradation often leads to a loss of potency.

A working knowledge of the functional groups within drug molecules, those that are likely to be susceptible to metabolism and those more likely to be resistant should be useful when counseling patients on the storage and use of their prescribed medicines. A common example is dispensing medicines into monitored dosage systems (MDS), where medicines are removed from manufacturers' packaging and placed in an environment where stability is unknown. Similarly, it is useful to have knowledge of functional groups whenever alternative therapeutic agents within a given chemical class are being considered as part

of a medicines use review.

2. Chemical Reactions

Most drugs are small organic molecules. The only differences between one drug and another are the number and type of functional groups present, and the connectivity between these groups. This is true of all organic molecules and, as such, there is little difference between drugs and other small organic molecules, such as those found in food and household products. The exception is that drugs have been specifically designed and optimized (over many years and costing hundreds of millions of pounds)¹ to have the desired pharmacological profile to enable them to treat a disease. With this in mind, it should therefore not be surprising that drugs undergo chemical reactions in much the same way as other organic molecules under set conditions. The main chemical reactions that affect the stability of a drug are oxidation and hydrolysis. Oxidation involves the removal of electrons from a molecule (or the addition of oxygen) and such reactions can be initiated by light, heat or certain trace metals.¹

Despite oxidation being a relatively common pathway for drug decomposition, it has not been studied in as

* Corresponding author.

E-mail address: rajdanish2k@gmail.com (Senthilraja. M).

much detail as hydrolysis since oxidative degradation can often be reduced to acceptable levels by storing susceptible drugs in the absence of light example use of amber vials and oxygen example store under nitrogen or argon or by using antioxidants in the formulation. Hydrolysis is the more common pathway for drug breakdown and is the mechanism that we will concentrate on here with regard to both chemical stability and metabolism.

3. Hydrolysis

Hydrolysis means the reaction of a molecule with water resulting in the cleavage of a chemical bond within that molecule. Although there are a large number of functional groups that are susceptible to hydrolysis, esters and amides are the most common ones found in drugs prone to hydrolysis. Figure 1 explains that the mechanism that accounts for the reaction of an ester ($X=OR$) or an amide ($X=NR_1R_2$, where R, R₁ and R₂ can be any arbitrary structure) resulting in the cleaved reaction products of a carboxylic acid and either an alcohol ($XH=ROH$) or an amine ($XH=R_1R_2NH$), respectively. Oxygen can draw electrons towards itself more strongly than carbon, so the C=O double bond in the ester or amide becomes polarized, leaving the oxygen slightly negatively charged and the carbon slightly positively charged.

As a result of this polarization, the electrons on water's oxygen atom are attracted to the slightly positive charge of the aforementioned carbon atom (unlike charges attract one another), resulting in hydrolysis (Figure 1, indicated by the bottom arrow at 2). This polarization and subsequent interaction is enhanced if the carbonyl oxygen is protonated (as shown at 1 in Figure 1).^{2,3}




Fig. 1: Simplified general hydrolysis mechanism of esters and amides

The reaction of water with esters proceeds more rapidly than with amides, and these reactions can be catalyzed by both acid and base. The rate is dependent on the pH of the aqueous solution. Similarly, the in vivo metabolism of esters and amides in drugs via a hydrolysis mechanism is catalyzed by hydrolytic enzymes present in various tissues and in plasma.

The reason for the difference in the rates of ester and amide hydrolysis lies in the structural difference between these two functional groups. The ester contains an oxygen atom, whereas the amide possesses a nitrogen atom at the

same position. This difference results in the carbon atom of the carbonyl group (C=O) being even more positively charged in the ester than in the amide and, as such, there is a greater attraction between that carbon and a water molecule. The opposite is true for the amide because the presence of the nitrogen atom in amides lessens the positive charge on the carbonyl carbon atom and thus it is less able to attract incoming water molecules.

This difference in the hydrolysis of esters and amides is exemplified with lidocaine and procaine (Figure 2). The ester-containing local anesthetic procaine is now seldom used.⁴⁻⁶ Although it is a good local anesthetic, its effect is short-lasting because its ester group is rapidly hydrolyzed. Conversely, lidocaine contains an amide bond, which is less readily hydrolyzed than procaine's ester. As such, it is more stable to the effects of hydrolysis and this, in addition to lidocaine's more bulky structure, makes it a longer-acting local anaesthetic.




Fig. 2: Chemical structures of lidocaine and procaine

Furthermore, the attention-deficit hyperactivity disorder drug methylphenidate (Figure 3) contains a methyl ester that is readily hydrolyzed by hydrolytic enzymes to yield ritalinic acid as the major (inactive) urinary metabolite in humans.⁷ Such a readily hydrolysable functional group poses a problem since it renders the drug inactive and, therefore, the drug needs to be administered more than once daily to maintain therapeutic levels. As such, a great deal of research into modified-release formulations has resulted in once-daily Concerta XL and Equasym XL as a means to address the non-compliance associated with self-administration during school hours. On the other hand, hydrolysis can also be helpful. The anti-motility drug co-phenotrope contains the ester diphenoxylate (Figure 3), which is readily hydrolyzed in humans to the related carboxylic acid, diphenoxylate, a five-times more potent antidiarrheal agent than the parent ester.

4. Prodrug S strategy

Salicylic acid is a good painkiller. However, the presence of the bare alcohol moiety causes gastric bleeding. Such an effect is overcome by masking this alcohol group as an ester — giving us aspirin — which is later hydrolyzed in the body to free the active drug. Such an approach is known as a prodrug strategy. The hydrolysis of aspirin also produces acetic acid as a product. In the poor storage of aspirin tablets, a smell of vinegar that pervades




Fig. 3: Chemical structures of methylphenidate and diphenoxylate

over time is a sign that hydrolysis is occurring. Another prodrug example includes the angiotensin-converting enzyme inhibitor enalapril, whose ester is hydrolyzed in vivo to generate the active carboxylic acid derivative enalaprilate (enalaprilate being used as an intravenous form of enalapril for hypertensive emergencies). Type I statins, such as simvastatin, which contain a cyclic ester (a lactone), and are hydrolyzed by body enzymes to produce the ring-opened, pharmacologically active hydroxy-acid common to these cholesterol-lowering drugs.

5. Amide-Containing Drugs

As previously mentioned, amide-containing drugs are also susceptible to hydrolysis, but it occurs at a much slower rate than the hydrolysis of esters. An example is the hydrolysis of the strained cyclic-amide ring (amides that are part of a ring structure are known as lactams) of β -lactam antibiotics. It is routine to prepare aqueous pediatric suspensions of penicillin antibiotics immediately before dispensing them since they are insufficiently stable to be supplied and stored long-term in water. Such suspensions should also be kept in the refrigerator to minimize hydrolysis of the strained lactam ring. The β -lactam antibiotics are also susceptible to hydrolysis through the action of hydrolytic enzymes. Such chemical instability and its associated implications can have a direct effect on the chosen route of administration. Only one has to look at some of the compounds that are incompatible with continuous infusion (eg, penicillin's and cephalosporins), to understand why both of which contain a strained and hydrolysis-prone β -lactam ring.

6. Other Functional Groups

Esters and amides are the most commonly encountered groups in drugs that are susceptible to hydrolysis. However, a large number of other functional groups can also react with water, resulting in broken bonds. Examples of such functional groups and the drugs in which they occur are imines ($C=N$) found in diazepam; acetals ($C(OR)_2$) found in digoxin; sulphates ($ROSO_3^-$) found in heparin; and phosphate esters ($ROPO_3^{2-}$), found in hydrocortisone sodium phosphate.

7. Preventing Hydrolysis

Although hydrolysis, as outlined above, can be problematic, it can be prevented or minimized in a number of ways. For in vivo metabolism, the preventive measures are a little trickier to overcome. However, hydrolysis can be prevented by chemically modifying the structure of the active compound in the early drug development stage, providing that the problematic hydrolysis is identified early enough. However, for inherently unstable drugs, preventive measures can include altering the dosage form in a number of ways. For liquid dosage forms, the hydrolysis of drugs is dependent on the presence of water, and thus storage of dry powders followed by reconstitution in water before dispensing minimizes hydrolysis.

Similarly, if a drug is known to be susceptible to hydrolysis at room temperature, its storage in a cool place is advised, and patients can be counseled on this practice. Additionally, pharmacists should provide the correct labeling on the packaging.

Moreover, the rate of hydrolysis is temperature-dependent and thus heat sterilization of such drugs could be a problem. For semi-solid dosage forms (ointment and creams), the stability of the active ingredient can be controlled by changing the nature of the ointment or cream base in the formulation. The sensitivity of a drug to hydrolysis in a solid dosage form can be controlled in a similar manner by preparing a less hygroscopic salt of the drug or by reducing the water content in the excipients used in the formulation.

8. Conclusion

Pharmacists responsibility to provide a safe and effective product for patients requires knowledge of the basic chemistry of drug stability. Common applications in most pharmacies are consideration of drug stability in MDSs. Although definitive stability data are often lacking, guidance is available on medicines that can safely be dispensed into MDSs. Knowledge of drug chemistry will assist pharmacists interpretation of this leading to improved patient safety.

9. Conflict of Interest

The authors declare no relevant conflicts of interest.

10. Source of Funding

None.

References

1. Snape TJ, Astles AM, Davies J. Understanding the chemical basis of drug stability and degradation. *Pharm J.* 2010;285(7622):410–6.
2. Reynolds DW, Facchine KL, Mullaney J. Available guidance and best practices for conducting forced degradation studies *Pharm. Technol.* 2002;26(2):48–56.

3. Brummer H. How to approach a forced degradation study. *Life Sci Technol Bull.* 2011;31:1–4. Available from: <https://www.sgs.com/~media/Global/Documents/Technical%20Documents/SGS-LSS-Forced%20Degradation-EN-11.pdf>.
4. Kats M. Forced degradation studies: regulatory considerations and implementation. *BioPharm Int.* 2005;18(7):40–5.
5. Maheswaran R. FDA perspectives: scientific considerations of forced degradation studies in ANDA submissions *Pharm. Technol.* 2012;36(5):73–80.
6. Klick S, Pim GM, Waterval J. Toward a generic approach for stress testing of drug substances and drug products. *Pharm Technol.* 2005;29(2):48–66.
7. Church C, Smith J. *the Pharmaceutical Journal.* 2006;276:75–79.

Author biography

Senthilraja. M, Professor & Principal

Cite this article: Senthilraja. M. A chemical rationale of drug stability and degradation- An insightful approach. *IP Int J Comprehensive Adv Pharmacol* 2022;7(2):102-105.