



Original Research Article

Acceptance or rejection of the COVID-19 Vaccine: A cross-sectional study among Orthodontists

Zainab Mousa Kadhom¹, Alaa Faleh Albo Hassan¹, Suha Ali Abdul Ameer²,
Mohammed Nahidh¹, Maria Maddalena Marrapodi³, Marco Cicciù⁴,
Giuseppe Minervini^{5,6*}

¹Dept. of Orthodontics, College of Dentistry, University of Baghdad, Baghdad, Iraq

²Ashur University, Baghdad, Iraq

³Dept. of Woman, Child and General and Specialist Surgery, University of Campania "Luigi Vanvitelli", Naples, Italy

⁴Dept. of Biomedical and Surgical and Biomedical Sciences, Catania University, Catania, Italy

⁵Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Tamil Nadu, Chennai, India

⁶Multidisciplinary Department of Medical-Surgical and Odontostomatological Specialties, University of Campania "Luigi Vanvitelli", Naples, Italy



ARTICLE INFO

Article history:

Received 06-05-2024

Accepted 15-05-2024

Available online 08-02-2025

Keywords:

COVID19

vaccine

orthodontists

health

ABSTRACT

Background: Close contact with the infected patient is one reason for the spread of infection. COVID-19 can easily infect orthodontists; it may be found in saliva droplets during the removal or insertion of the orthodontic fixed appliance components or even during the removal of the removable appliance.

Aims: This study aims to assess the acceptance or rejection of the COVID-19 vaccine among Iraqi orthodontists.

Materials and Methods: A web-based cross-sectional survey among orthodontists in Iraq was conducted. Thirteen pre-structured questions were sent in a Google Form via their WhatsApp and Telegram groups.

Results: There were 123 orthodontists participating in this survey. The majority of the participants (96.75%) were aware of COVID-19, and 62.60% suggested that coronavirus poses a significant risk. Orthodontists who took the vaccine accounted for 93.50%; among them, those who preferred the Pfizer vaccine accounted for 57.39%. The side effects associated with the vaccine were the leading reasons for vaccine refusal.

Conclusions: Although there is a high level of awareness about COVID-19 viruses among orthodontists, some abstained from the vaccine because of fear. The acceptance of COVID-19 vaccines and the number of participants who got the vaccine were high to achieve precise infection control in orthodontic clinics and decrease viral infection spread.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial 4.0 International](https://creativecommons.org/licenses/by-nc/4.0/), which allows others to remix, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The COVID-19 outbreak originated in Wuhan, China, in December 2019, leading to significant disruptions in

healthcare systems globally. Severe viral pneumonia leads to acute respiratory syndrome with clinical symptoms of infection that are not always specific.^{1,2}

Healthcare workers are at high risk of infection because they are directly involved in diagnosing, treating, and caring for patients.³ Dentistry and its related internships are one of the most deeply influenced sectors. Dental professionals are

* Corresponding author.

E-mail address: giuseppe.minervini@unicampania.it (G. Minervini).

exposed to high-risk environments due to the presence of oral fluids and aerosols, increasing the likelihood of viral exposure and transmission.^{4,5}

The infection spreads mainly through respiratory droplets and during close contact; symptomatic, pre-symptomatic, and asymptomatic patients can carry it.

WHO governments attempted to contain COVID-19 because of its highly infectious nature. Antiviral medications and immunotherapy (monoclonal antibodies and protease inhibitors) are potential therapeutics being developed or in different stages of clinical trials.^{6,7}

The vaccine is the most effective and frequently used way to prevent disease and control infection.⁸ Therefore, vaccination may be the most reliable way to stop and contain COVID-19.⁹ COVID-19 vaccine development and supply are an ongoing process.⁸ In North America and Europe, well-known companies released many candidate vaccines for high-risk populations and healthcare workers, such as older adults and patients with chronic diseases.³

There are 138 vaccine candidates; their mode of action depends on the immune response to its integral parts (DNA, RNA, or protein), and about 21 of these vaccines are approved worldwide for emergency use.⁹ Pfizer-BioTech, Moderna, and Johnson & Johnson developed three vaccines that are being administered broadly.¹⁰

The Pfizer-BioNTech BNT162b2 vaccine was approved by the FDA on August 23, 2021, after meeting the FDA requirements for manufacturing, efficacy, and safety.¹¹ Other approved vaccines are BBIBP, Sinopharm-Wuhan (China), CoronaVac, CoviVac (Russia), QazCovid-in (Kazakhstan), and COVIran Barekat (Iran).⁹

There has been a lot of research on the knowledge, attitude, and behaviors during COVID-19 pandemic in addition to the vaccine acceptance among orthodontists, general dentists and health coworkers.^{12–21} The present study is the first Iraqi study, and it aims to assess the acceptance and rejection of the COVID-19 vaccine among Iraqi orthodontists.

2. Materials and Methods

The College of Dentistry, University of Baghdad's ethical committee approved this cross-sectional survey. The sample size was estimated using the online SurveyMonkey sample size calculator; the population size was 180, representing the current total number of orthodontists in Iraq, with a confidence interval of 95% and a margin of error of 5; hence, the required sample size was 123.

The survey included 13 questions regarding COVID-19 vaccines. It was disseminated by Google Forms and sent as a link through the participants' WhatsApp and Telegram groups. This survey was anonymous to ensure the information's confidentiality.

The questionnaire had two parts. The first collected demographic data such as gender, age, qualification, and

place of work. The second part was concerned with questions related to the COVID-19 vaccine. Microsoft Office Excel analyzed the collected responses as frequencies and percentages.

3. Results

This study's results are based on the responses of 123 of 180 Iraqi orthodontists invited to participate in this online questionnaire regarding their acceptance or rejection of the COVID-19 vaccine.

Table 2 shows the demographic information. Half of the respondents were aged 36–45, 83.74% had a master's degree, and 82.11% worked in both institutional and private clinics.

In Table 2, 96.75% of participants were aware of COVID-19, and 54.47% were infected along with their families. Moreover, 62.60% of participants thought that the coronavirus poses a major risk; 47.15% agreed that the COVID-19 vaccine effectively protects from COVID-19. Furthermore, 91.87% of the respondents encouraged their patients to take the vaccine; 93.5% of the participants took the vaccine, and 6.5% did not because of fear of the side effects that could occur after taking the vaccine and the absence of scientific information about it.

Table 3 shows the vaccinated participants' responses. The Pfizer vaccine was preferred by 57.39%. The orthodontists who changed their infection control protocols after vaccination accounted for 54.78%; and 68.70% did not catch the virus after taking the vaccine compared to 9.57% who caught it. Moderate symptoms of infection were found in 45.45% of all orthodontists. Finally, 81.82% of the participants brought antivirus protection to work.

4. Discussion

The role of dentists and dental students in the fight against COVID-19 is of paramount value²² in terms of providing safe oral/dental care during the pandemic and educating and motivating the public to accept COVID-19 vaccines and adhere to preventive measures.¹⁸

A survey taken in Iraq during January 2021 among the general population suggested that only 77.6% of Iraqis were willing to get the vaccine as soon as it was available.²³ The current survey was conducted on orthodontists since all dentists, including orthodontists, can be infected by the virus;⁷ however, orthodontic practice produces fewer aerosols than other dental specialties. Clear aligners, attachment bonding, bracket insertion, and residual adhesive removal create aerosol; COVID-19 may also be found in saliva droplets, which might splash while removing or inserting orthodontic ligatures and wires.²⁴

Most guidelines do not contain sufficient information about orthodontic management during COVID-19. Because orthodontic treatment is continuous and needs follow-up

Table 1: Demographic data of the participants

Parameters		N	%
Age	25-35	48	39.02
	36-45	62	50.41
	46-55	13	10.57
	Total	123	100
Genders	Male	62	50.41
	Female	61	49.59
	Total	123	100
Qualification	Certificate	5	4.07
	Master	103	83.74
	Ph.D.	15	12.20
	Total	123	100
Place of work	Private	8	6.50
	Institutional	14	11.38
	Both	101	82.11
	Total	123	100

Table 2: Awareness to COVID-19 and vaccines

Questions	Responses	N	%
Are you aware of COVID-19?	Yes	119	96.75
	No	4	3.25
	Total	123	100
Have you or any members of your family been infected with COVID-19 in the past?	I and my family were	67	54.47
	I don't sure	8	6.50
	I was only	5	4.07
	No one has been infected	14	11.38
	Someone in the family only	29	23.58
To which extent do you think coronavirus poses a risk to you personally?	Total	123	100
	Major	77	62.60
	Moderate	45	36.59
	No risk	1	0.81
	Total	123	100
Does COVID-19 vaccine effective in protecting you from COVID-19?	Strongly agree	32	26.02
	Agree	58	47.15
	Neutral	32	26.02
	Disagree	1	0.81
Do you encourage your patients to take the vaccine?	Total	123	100
	Yes	113	91.87
	No	2	1.63
	I do not care	8	6.50
Did you take the vaccine?	Total	123	100
	Yes	115	93.50
	No	8	6.50
	Total	123	100
Why don't you want to take the vaccine?	I am afraid of the side effects that may occur after taking the vaccine	5	62.5
	I do not have enough scientific information about it because it is new	2	25
	The vaccine was not available	1	12.5
	I trust my own immunity to protect me	0	0
	Someone else told me that the vaccine is not safe	0	0
	Total	8	100

Table 3: Responses of vaccinated participants about vaccines

Questions	Responses	N	%
Which of the following types of vaccine do you prefer?	Chinese	26	22.61
	Oxford (Astra Zeneca)	21	18.26
	Pfizer	66	57.39
	US Moderna	2	1.74
	Total	115	100
Did your application of health protection instructions against the corona virus change after taking the vaccine?	Yes	63	54.78
	No	52	45.22
	Total	115	100
	Yes	11	9.57
Did you catch the corona virus after taking the vaccine?	No	79	68.70
	I am not sure	25	21.74
	Total	115	100
	Mild	4	36.36
How were the symptoms of infection?	Moderate	5	45.45
	Severe	2	18.18
	Total	11	100
	Fever	1	9.09
	Fever, weakness, and fatigue	2	18.18
What symptoms did you have after taking a first and a second dose of the vaccine?	Flu-like symptoms	3	27.27
	Weakness and fatigue	1	9.09
	Weakness, fatigue, flu-like symptoms	1	9.09
	All of the above	3	27.27
	Total	11	100
	I go to the clinic with protection means	9	81.82
How did you deal with your patients next day after the first and second vaccine doses?	I go to treat only the urgent cases	1	9.09
	I did not go to the clinic	1	9.09
	Total	11	100

every 4–6 weeks, true orthodontic emergencies during a crisis like this should be clarified, and protocols should be developed on how to manage them.¹³

In this study, orthodontists showed a high level of awareness about COVID-19 viruses, as did orthodontists in China¹³ and Turkey¹⁴, but another study¹⁵ showed a moderate knowledge of COVID-19 vaccination.

In Iurcove et al.'s study,¹⁹ which was conducted on dentists, higher infection rates among family members were recorded than in the present study. So, during the lockdown and before the vaccine was available, some orthodontists depended on social media to communicate with and supervise their patients from afar.²⁵ Social distancing and staying away from family are preventive measures that must be taken during a pandemic.

About 62.60% of the sample thought the coronavirus poses a major risk. This supports taking the preventive measures recommended in dentistry by encouraging vaccination, which is now widespread. The hesitation in taking it and the misleading information and rumors make it a significant challenge because dentists are at high risk of infection.²⁶ Other potential factors that could influence vaccine acceptance are risk awareness and safety perception

among dental professionals. Dentists have been shown to have an acceptable level of risk awareness.²⁷

The vaccine's effectiveness showed geographical variations; in the USA, the efficacy was 72%, but in this study, only 47.15% of the orthodontists thought the COVID-19 vaccine effectively protected from the coronavirus. As in the current study, many orthodontists encouraged their patients to get the vaccine to protect them from infection. The USA prioritized the vaccine for American dental and medical students,²⁸ so giving priority to dentists to receive the COVID vaccine is ethical and necessary to ensure the community's health. Patients may feel reassured when visiting the dentist if they know that a high percentage of their dentist and dental staff were vaccinated and encouraged their patients to get the vaccine.²⁰

In many studies,^{12,29} a high number of participants were vaccinated, as in the present study, and the most preferred vaccine was Pfizer's vaccine rather than the others^{15,29–31}, since mRNA vaccines are an excellent alternative to support immunity against viruses.³⁰ The most common reason for vaccine refusal among the healthcare workers and staff at Ugandan³¹ and Nepalese³² medical colleges was concern

about the vaccine's safety. The side effects that may occur after taking the vaccine were the primary cause of fear of the vaccine in the present study.

After vaccination, side effects typically comprised a small amount of fatigue with a headache, elevated body temperature, and soreness at the injection site; sometimes, the symptoms were mild^{15,29,33,34}. In addition, our study noted flu-like symptoms. Some respondents believed that COVID-19 would go away permanently when they changed their ways of practicing the profession; orthodontists and doctors indicated that they would improve infection control methods.¹⁶ N95 respirators and face shields became the norm rather than the exception, particularly for aerosol-generating procedures. Most practices have also begun to adhere to occupational safety and health administration requirements to test the fit of their N95 respirators at least annually, in addition to developing and training team members in their own respiratory protection programs.³⁵

Orthodontic clinics must follow thorough and precise infection control measures to meet patients' needs. To minimize the spread of viral infection, all precautionary measures for COVID-19 have to be made available in the clinic, in addition to continuous dental healthcare training for workers and patients.^{36–41}

Vaishya et al.²¹ determined that 2.63% of vaccinated healthcare coworkers acquired the SARS-CoV-2 infection after vaccination; this percentage is less than the 9.57% of our sample who were infected after vaccine uptake.

Some orthodontists in this study changed their health protection protocols against the coronavirus after taking the vaccine to prevent the infection after immunization, so a high percentage of the sample did not catch the coronavirus after taking the vaccine. This supports the Pietrzak and Hanke¹⁷ study, which stated that healthcare workers were four times less likely to develop asymptomatic COVID-19 infection 12 days after receiving the vaccine.

The study's limitation was that the sample included only orthodontists and did not contain all specializations in dentistry.

5. Conclusions

Although orthodontists are highly aware of COVID-19 viruses, some refuse the vaccine out of fear. The acceptance of COVID-19 vaccines and the number of participants who took the vaccine were high.

6. Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

7. Source of Funding

This study did not receive any funding in any form.

8. Conflict of Interest

None.

References

- Adham ZS, Bh AG. Prevalence of viral co-infection among COVID-19 cases in association with disease severity and oral hygiene. *J Bagh Coll Dent*. 2021;33(3):1–8.
- Hussein SI, Abd ST, Al-Khayat FA, Mahmood HK. Complete blood count and saliva parameters as an indicator for infected patients with coronavirus COVID-19. *J Bagh Coll Dent*. 2023;35(1):76–85.
- Ali S, Noreen S, Farooq I, Bugshan A, Vohra F. Risk assessment of healthcare workers at the frontline against COVID-19. *Pak J Med Sci*. 2020;36(4):99–103.
- Bentley CD, Burkhart NW, Crawford JJ. Evaluating spatter and aerosol contamination during dental procedures. *J Am Dent Assoc*. 1994;125(5):579–84.
- Checchi V, Bellini P, Bencivenni D, Consolo U. COVID-19 dentistry-related aspects: a literature overview. *Int Dent J*. 2020;71(1):21–6.
- Aminjafari A, Ghasemi S. The possible of immunotherapy for COVID-19: A systematic review. *Int Immunopharmacol*. 2020;83:106455.
- Wang J, Peng Y, Xu H, Cui Z, Williams RO. The COVID-19 vaccine race: Challenges and opportunities in vaccine formulation. *AAPS Pharm Sci Tech*. 2020;21(6):225.
- Lurie N, Saville M, Hatchett R, Halton J. Developing Covid-19 vaccines at pandemic speed. *N Engl J Med*. 2020;382(21):1969–73.
- Covid-19 Tracker. COVID 19 vaccine tracker. Available from: <https://covid19.trackvaccines.org>.
- Timeline: WHO's COVID-19 response. World Health Organization; 2021. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/interactive-timeline>.
- FDA approves first COVID-19 vaccine; 2021. Available from: <https://www.fda.gov/news-events/press-announcements/fda-approves-first-covid-19-vaccine>.
- Al-Azzawi M, Alghazali MW, Hareeja MM, Abbas IK, Alwan AH. COVID-19 vaccine hesitancy among Iraqi dentists. *J Med Chem Sci*. 2023;6(2):424–33.
- Hua F, Qin D, Yan J, Zhao T, He H. COVID-19 related experience, knowledge, attitude, and behaviors among 2,669 orthodontists, orthodontic residents, and nurses in China: A cross-sectional survey. *Front Med*. 2020;7:481.
- Yilmaz HN, Ozbilen EO. The assessment of knowledge, behaviors, and anxiety levels of the orthodontists about COVID-19 pandemic. *Turk J Orthod*. 2020;33(4):224–31.
- Nasr L, Saleh N, Hleyhel M, El-Outa A, Noujeim Z. Acceptance of COVID-19 vaccination and its determinants among Lebanese dentists: A cross-sectional study. *BMC Oral Health*. 2021;21(1):484.
- Isiekwe IG, Adeyemi TE, Aikins FA, Umeh OD. Perceived impact of the COVID-19 pandemic on orthodontic practice by orthodontists and orthodontic residents in Nigeria. *J World Fed Orthod*. 2020;9(3):123–8.
- Pietrzak P, Hanke W. COVID-19 and dentistry-safety issues regarding doctor and patient situation in time of vaccine availability. *Med Pr*. 2021;72(6):729–37.
- Nassani MZ, Noushad M, Rastam S. Determinants of COVID-19 vaccine acceptance among dental professionals: A multi-country survey. *Vaccines (Basel)*. 2022;10(10):1614.
- Iurcov R, Pop LM, Ciavoi G, Iorga M. Evaluating the practice of preventive behaviors and the fear of COVID-19 among dentists in Oradea metropolitan area after the first wave of pandemic; a cross-sectional study. *Healthcare (Basel)*. 2021;9(4):443.
- Wilson R, Jonke G. The ethics of dentists receiving the COVID-19 vaccine. *J Am Dent Assoc*. 2021;152(5):408–9.
- Vaishya R, Sibal A, Malani AK, Prasad H. SARS-CoV-2 infection after COVID-19 immunization in healthcare workers: A retrospective, pilot study. *Indian J Med Res*. 2021;153(5-6):550–4.

22. Crowder L. Dental professionals' role in the fight against COVID-19: Current evidence. *Evid Based Dent.* 2021;22(2):62–3.
23. Ghazi HF, Taher TMJ, Alfadhul S. Acceptance of COVID-19 vaccine among general population in Iraq. *Iraqi National J Med.* 2021;3(1):93–103.
24. Silva RS, Jardim ACG, Siqueira WL. Coronavirus COVID-19 impacts to dentistry and potential salivary diagnosis. *Clin Oral Investig.* 2020;24(4):1619–21.
25. Nahidh M, Al-Khawaja NFK, Jasim HMJ, Cervino G, Cicciù M, Minervini G, et al. The role of social media in communication and learning at the time of COVID-19 lockdown-an online survey. *Dent J (Basel).* 2023;11(2):48.
26. Fontanet A, Autran B, Lina B, Kieny MP, Karim SSA, Sridhar D. SARS-CoV-2 variants and ending the COVID-19 pandemic. *Lancet.* 2021;397(10278):952–4.
27. Wolf TG, de Col L, Rad SB, Castiglia P, Arghittu A, Cannavale M, et al. How the COVID-19 pandemic affects risk awareness in dentists: A scoping review. *Int J Environ Res Public Health.* 2022;19(9):4971.
28. Kelekar AK, Lucia VC, Afonso NM, Mascarenhas AK. COVID-19 vaccine acceptance and hesitancy among dental and medical students. *J Am Dent Assoc.* 2021;152(8):596–603.
29. Jękowiak I, Wiatrak B, Dziewiszek PG, Szeląg A. The incidence and severity of post-vaccination reactions after vaccination against COVID-19. *Vaccines.* 2021;9(5):502.
30. Xu S, Yang K, Li R, Zhang L. mRNA vaccine era-mechanisms, drug platform and clinical prospect. *Int J Mol Sci.* 2020;21(18):6582.
31. Kanyike AM, Olum R, Kajjimu J. Acceptance of the coronavirus disease- 2019 vaccine among medical students in Uganda. *Trop Med Health.* 2021;49(1):37.
32. Paudel S, Palaian S, Shankar PR, Subedi N. Risk perception and hesitancy toward covid-19 vaccination among healthcare workers and staff at a medical college in Nepal. *Risk Manag Healthc Policy.* 2021;14:2253–61.
33. Almuftu HB, Mohammed SA, Abdullah AM, Merza MA. Potential adverse effects of COVID19 vaccines among Iraqi population; a comparison between the three available vaccines in Iraq; a retrospective cross-sectional study. *Diab Metab Syndr.* 2021;15(5):102207.
34. Sa'ady A, Abdulrasol AT, Obaid ZA, Alhindy AF, Al-Mumin H. Prevalence of adverse effects from COVID-19 vaccine among Iraqi adults: A retrospective cross-sectional study. *J Emerg Med Trauma Acute Care.* 2022;2022(3):1–6.
35. Dentistry Worker and Employers Occupational Safety and Health Administration; 2021. Available from: <https://www.osha.gov/coronavirus/control-Prevention/Dentistry>.
36. Rafeeq RA, Saleem AE, Nahidh M, Kadhum AS, Al-Huwaizi AF, Marrapodi MM, et al. Clinical management and infection control protocols during the COVID-19 pandemic: An online survey. *Technol Health Care.* 2023;31(5):1579–92.
37. Venugopal A, Vaid N, Bowman SJ. Outstanding, yet redundant? After all, you may be another Choluteca Bridge. *Semin Orthod.* 2021;27(1):53–6.
38. Lee YJ, Kook YA, Park JH, Park J, Bayome M, Vaid NR. Short-term cone-beam computed tomography evaluation of maxillary third molar changes after total arch distalization in adolescents. *Am J Orthod Dentofac Orthop.* 2019;155(2):191–8.
39. Vaid N, Doshi V, Vandekar M. What's "Trend"ing in Orthodontic literature? *APOS Trends Orthod.* 2016;6(1):1.
40. Awad MG, Ellouze S, Ashley S, Vaid N, Makki L, Ferguson DJ, et al. Accuracy of digital predictions with CAD/CAM labial and lingual appliances: A retrospective cohort study. *Semin Orthod.* 2018;24(4):393–406.
41. Alansari R, Vaid N. Why do patients transition between orthodontic appliances? A qualitative analysis of patient decision-making. *Orthod Craniofac Res.* 2023;27(3):439–46.

Author's biography

Zainab Mousa Kadhom, Dental Surgeon

Alaa Faleh Albo Hassan, Dental Surgeon

Suha Ali Abdul Ameer, Dental Surgeon

Mohammed Nahidh, Dental Surgeon

Maria Maddalena Marrapodi, PhD Scholar

Marco Cicciù, Dental Surgeon

Giuseppe Minervini, -

Cite this article: Kadhom ZM, Faleh Albo Hassan A, Ali Abdul Ameer S, Nahidh M, Maddalena Marrapodi M, Cicciù M, Minervini G. Acceptance or rejection of the COVID-19 Vaccine: A cross-sectional study among Orthodontists. *J Contemp Orthod* 2025;9(1):48-53.